

**Momentum Spectra in the
Current Region of the Breit Frame
in Deep Inelastic Scattering
at HERA**

N. Brook

on behalf of the ZEUS collaboration

Abstract

The production of charged particles has been measured in deep inelastic scattering (DIS) with the ZEUS detector. The evolution of the moments of the scaled momenta distributions in Q^2 and x has been investigated in the current fragmentation region of the Breit frame. The results in the current region are compared to e^+e^- data and QCD analytical calculations. The results are consistent with the universality of single-particle spectra in DIS and e^+e^- annihilation at high Q^2 . The results at low Q^2 disagree with analytical calculations based on the modified leading log approximation (MLLA) and local parton hadron duality (LPHD).

*talk given at the
3rd UK Phenomenology Workshop
on HERA Physics,
Durham, United Kingdom.
Sept 20-25, 1998*

Introduction

This paper reports the results of a study of the properties of the hadronic final state in positron-proton deep inelastic scattering (DIS). The event kinematics of DIS are determined by the negative square of the four-momentum transfer of the virtual exchanged boson, $Q^2 \equiv -q^2$, and the Bjorken scaling variable, $x = Q^2/2P \cdot q$, where P is the four-momentum of the proton. In the Quark Parton Model (QPM), the interacting quark from the proton carries four-momentum xP . The variable y , the fractional energy transfer to the proton in its rest frame, is related to x and Q^2 by $y \approx Q^2/xs$, where $\sqrt{s}$ is the positron-proton centre of mass energy.

A natural frame in which to study the dynamics of the hadronic final state in DIS is the Breit frame [1]. In this frame the exchanged virtual boson is completely space-like and has a four-momentum $q = (0, 0, 0, -Q = -2xP^{Breit}) \equiv (E, p_x, p_y, p_z)$, where P^{Breit} is the momentum of the proton in the Breit frame. The particles produced in the interaction can be assigned to one of two regions: the current region if their z -momentum in the Breit frame is negative, and the target region if their z -momentum is positive. The advantage of this frame is that it gives a maximal separation of the incoming and outgoing partons in the QPM. In this model the maximum momentum a particle can have in the current region is $Q/2$.

The current region in the Breit frame is analogous to a single hemisphere of e^+e^- annihilation. In $e^+e^- \rightarrow q\bar{q}$ annihilation the two quarks are produced with equal and opposite momenta, $\pm\sqrt{s}/2$. The fragmentation of these quarks can be compared with that of the quark struck from the proton which has outgoing momentum $-Q/2$ in the Breit frame. In the direction of this struck quark the scaled momentum spectra of the particles, expressed in terms of $x_p = 2p^{Breit}/Q$, are expected to have a dependence on Q similar to that observed in e^+e^- annihilation [2, 3, 4] at energy $\sqrt{s} = Q$.

Within the modified leading log approximation (MLLA) there are predictions of how the higher order moments of the parton momentum spectra should evolve with energy scale [5, 6]. The parton level predictions depend on two free parameters, a running strong coupling, governed by a QCD scale Λ , and an energy cut-off, Q_0 , below which the parton evolution is truncated. The hypothesis of local parton hadron duality (LPHD) [7], which relates the observed hadron distributions to the calculated parton distributions via a constant of proportionality, is used in conjunction with the predictions of the MLLA allowing the calculation to be directly compared with data.

Results

The moments of the $\ln(1/x_p)$ distributions have been investigated up to the 4th order; the mean (l), width (w), skewness (s) and kurtosis (k) were extracted from the distribution by fitting a distorted Gaussian of the following form:

$$\frac{1}{\sigma_{tot}} \frac{d\sigma}{d\ln(1/x_p)} \propto \exp\left(\frac{1}{8}k - \frac{1}{2}s\delta - \frac{1}{4}(2+k)\delta^2 + \frac{1}{6}s\delta^3 + \frac{1}{24}k\delta^4\right)$$

where $\delta = (\ln(1/x_p) - l)/w$, over a range of 3 units ($Q^2 < 160 \text{ GeV}^2$) or 4 ($Q^2 \geq 160 \text{ GeV}^2$) units in $\ln(1/x_p)$ around the mean. The equation was motivated by the expression used for the MLLA predictions of the spectra in ref. [5].

Figure 1 shows the skewness of the $\ln(1/x_p)$ spectra as a function of $\ln(Q)$. It is evident that the skewness decreases with increasing Q . Similar fits performed on e^+e^- data shows a reasonable agreement with our results at high Q^2 , consistent with the universality of fragmentation for this distribution. The ARIADNE Monte Carlo model [8] gives a reasonable description of the data. The data are compared with the MLLA predictions of ref. [6], using a value of $\Lambda = 175 \text{ MeV}$, for different values of Q_0 . The MLLA calculations predict a negative skewness which decreases towards zero with increasing Q in the case of the limiting spectra ($Q_0 = \Lambda$). This is contrary to the measurements. A reasonable description of the behaviour of the skewness with Q can be achieved for a truncated cascade ($Q_0 > \Lambda$), but a consistent description of the mean, width, skewness and kurtosis cannot be achieved [9]. A range of Λ values were investigated and none gave a good description of all the moments. We conclude that the MLLA predictions, assuming LPHD, do not describe the data. It should be noted though that a moments analysis has been performed [10], taking into account the limitations of the massless assumptions of the MLLA predictions, where good agreement was found between the limiting case of the MLLA and e^+e^- data over a large range of energy, $3.0 < \sqrt{s} < 133.0 \text{ GeV}$.




Figure 1: Evolution of the skewness of the $\ln(1/x_p)$ distribution with $\ln(Q)$. Data from e^+e^- and ep are shown together with, ARIADNE Monte Carlo predictions and the MLLA predictions of Dokshitzer *et al* [6] predictions (the full line is $Q_0 = \Lambda$, the dashed $Q_0 = 2\Lambda$, and the dotted $Q_0 = 3\Lambda$). The overlapping points are different x ranges in the same Q range. The inner error bars are the statistical error and the outer error bars are the systematic and statistical errors added in quadrature.

Summary

Charged particle distributions have been studied in the current region of the Breit frame in DIS. The moments of the $\ln(1/x_p)$ spectra in the current region at high Q^2 exhibit the same energy scale behaviour as that observed in e^+e^- data. The moments cannot be described by the MLLA calculations together with LPHD.

References

- [1] R. P. Feynman, ‘Photon-Hadron Interactions’, Benjamin, N.Y. (1972).
- [2] Yu. Dokshitzer et al., Rev. Mod. Phys. 60 (1988) 373.
- [3] A. V. Anisovich et al., Il Nuovo Cimento A106 (1993) 547.
- [4] K. Charchuła, J. Phys. G19 (1993) 1587.
- [5] C. P. Fong and B. R. Webber, Phys. Lett. B229 (1989) 289,
C. P. Fong and B. R. Webber, Nucl. Phys. B355 (1991) 54.
- [6] Yu. Dokshitzer, V. Khoze and S. Troyan, Int. J. Mod. Phys. A7 (1992) 1875.
- [7] Ya. Azimov et al., Z. Phys. C27 (1985) 65.
- [8] L.Lönnblad, ARIADNE version 4.03 program and manual, Comp. Phys. Comm. 71 (1992) 15.
- [9] ZEUS Collab., to be published.
- [10] S. Lupia and W. Ochs, Eur. Phys. J. C2 (1998) 307.